

Psychological Medicine

<http://journals.cambridge.org/PSM>


Additional services for *Psychological Medicine*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Different responses to K-10 and CIDI suggest that complex structured psychiatric interviews underestimate rates of mental disorder in old people

D. W. O'Connor and R. A. Parslow

Psychological Medicine / Volume 39 / Issue 09 / September 2009, pp 1527 - 1531

DOI: 10.1017/S0033291708004728, Published online: 02 December 2008

Link to this article: http://journals.cambridge.org/abstract_S0033291708004728

How to cite this article:

D. W. O'Connor and R. A. Parslow (2009). Different responses to K-10 and CIDI suggest that complex structured psychiatric interviews underestimate rates of mental disorder in old people. *Psychological Medicine*, 39, pp 1527-1531 doi:10.1017/S0033291708004728

Request Permissions : [Click here](#)

Different responses to K-10 and CIDI suggest that complex structured psychiatric interviews underestimate rates of mental disorder in old people

D. W. O'Connor^{1*} and R. A. Parslow²

¹ Department of Psychological Medicine, Monash University, Melbourne, Australia

² Australian Centre for Post-traumatic Mental Health, University of Melbourne, Melbourne, Australia

Background. Epidemiological surveys based on complex diagnostic interviews, such as the Composite International Diagnostic Interview (CIDI), report very low rates of anxiety and depressive disorders in older age groups. Mental health checklists show much less change over the lifespan. This paper explores the possibility that complex interviews present a special challenge to older respondents and thereby exaggerate the decline in mental disorder with age.

Method. Analysis of data from an Australian national mental health survey with 10 641 community-resident adult respondents. Measures of interest included ICD-10 anxiety and depression diagnoses, scores on the Kessler Psychological Distress Scale (K-10), agreement between K-10 and CIDI anxiety and depressive questions, and changes in agreement with age.

Results. Levels of inconsistency between simple and complex questions about anxiety and depression rose with age.

Conclusions. Older people may have difficulty attending to and processing lengthy, complex questionnaires. When in doubt, their preferred response may be to deny having experienced symptoms, thus deflating rates of diagnosed mental disorder. We recommend that simple mental health scales be included in epidemiological studies involving older age groups.

Received 31 March 2008; Revised 1 September 2008; Accepted 22 September 2008; First published online 2 December 2008

Key words: CIDI, diagnosis, K-10, mental disorder, old age, rating scales.

Introduction

A national survey commissioned in 1997 by the Australian Commonwealth Government reported that older, community-resident Australians were less prone than younger adults to anxiety disorders and depression (Henderson *et al.* 2000; O'Connor, 2006). Anxiety and depressive disorders both peaked in mid-life and then declined to such an extent that only 4.6% of people aged ≥ 65 years met ICD-10 criteria for an anxiety disorder in the previous 12 months and only 1.5% met criteria for major depression or dysthymia. By contrast, rates in the age band 45–54 years were 12.9% and 6.9% respectively.

Similar age-related falls in mental disorder have been reported in other countries (ESEMed/MHEDEA 2000 Investigators, 2004; Kessler *et al.* 2005; Wells *et al.* 2006; Lee *et al.* 2007) and persuasive arguments have

been marshalled by way of explanation. Older people might be less vulnerable to mental disorder because they were reared in tough times and cope better with adversity; their coping skills have been honed over decades, and they face fewer daily hassles (Henderson, 1994; Lindsay *et al.* 2006; Villamil *et al.* 2006; Yang, 2007).

It is striking, however, that most surveys reporting sharp falls in mental disorder with age have relied on lengthy, structured interviews such as the Diagnostic Interview Schedule (DIS), which generates DSM diagnoses (Robins *et al.* 1981), and the related Composite International Diagnostic Interview (CIDI), which generates both DSM and ICD diagnoses (WHO, 1994).

The CIDI is very complex indeed. In Version 1.20 used in Australia, the screening question for generalized anxiety disorder reads as follows: 'In the past 12 months, have you had a period of a month or more when for most of the time you felt worried, tense or anxious about everyday problems such as work or family?' A positive response prompted two supplementary questions: 'Has that period of feeling

* Address for correspondence: Professor D. W. O'Connor, Kingston Centre, Warrigal Road, Cheltenham, Victoria 3192, Australia.
(Email: Daniel.OConnor@med.monash.edu.au)

worried, tense or anxious been going on for six months or longer?' and 'In the past 12 months, did you have a time when you worried a lot more than most people would in your situation?' Positive responses to both these questions then prompted queries about 24 specific symptoms, their severity and persistence, and the content of worrisome thoughts. The process for depression follows a similar pathway.

Such questions require analysis of multiple symptoms, time-frames and attributions, and might therefore present a challenge to very old respondents. The ageing process, even in healthy older people, is associated with reduced attention span, working memory and information processing speed, all of which can impact on the capacity to respond quickly to batteries of multi-faceted questions (Anderson, 2008). In general, cognitive tasks that require the rapid manipulation of large quantities of complex new information become progressively more difficult with age. Physical frailty, pain, deafness and other age-related morbidities impose an additional burden.

Diagnostic interviews developed specifically for use with older people, including those with dementia, are typically less demanding. In the depression section of the Cambridge Examination for Mental Disorders of the Elderly (CAMDEX), for example, the question about low mood is simple: 'Do you feel sad or depressed or miserable?' (Roth *et al.* 1988). The Geriatric Mental State Schedule (GMS) takes a semi-structured approach with room for probes if required (Copeland *et al.* 1976). It is still much easier to complete than the CIDI and typically generates relatively high rates of 'depression syndrome cases'. In a small Australian study, 6.3% of the subjects aged 70–74 years who were assessed using the GMS met criteria for major depression, in contrast to 2.4% in the survey described here (Kay *et al.* 1985).

Psychiatric glossaries such as ICD-10 divide respondents into 'cases' of mental disorder and 'non-cases'. The results are clinically useful but somewhat arbitrary because 'caseness' can hinge on responses to a single question. Continuous ratings of mental health, by contrast, shed light on the whole community, including 'subthreshold cases', that is respondents who fail to meet criteria for diagnosis but might still be disabled and worthy of attention. Scales of mood and morale are typically brief and easy to complete with short time-frames, simple questions and small numbers of predetermined responses. They are therefore less subject to influence by age-related changes in cognition, sensory capacity and physical health.

It is of interest, therefore, to compare responses across the lifespan to simple and complex questions regarding mental health. The Australian national survey included two brief mental health scales, the

Kessler Psychological Distress Scale (K-10; Kessler *et al.* 2002) and the General Health Questionnaire (GHQ-12; Goldberg, 1972). We focused on K-10 in this study because its questions map better against CIDI.

Method

The Australian Bureau of Statistics surveyed a representative sample of residents of private dwellings throughout the country. Non-private dwellings such as hospitals and aged care facilities were excluded. One person aged ≥ 18 years from each of 13 600 eligible residences was invited to participate, of whom 78% agreed to be interviewed, giving a total of 10 641 respondents. All respondents completed the K-10, GHQ-12 and CIDI in that order. Respondents aged ≥ 65 years completed a cognitive scale, the Mini-Mental State Examination (MMSE; Folstein *et al.* 1975), and those who scored ≤ 18 points were not interviewed further.

Instruments

The CIDI anxiety and depression screening questions were put to all respondents and only those who endorsed them were questioned further. It is not possible, therefore, to compare the full suite of CIDI responses with other measures. The analyses below focus just on the screening questions for generalized anxiety disorder and major depression.

K-10 checks whether 10 mental symptoms have been present in the past 4 weeks 'for all, most, some, a little or none of the time'. Scores range from 10 to 50, with higher scores denoting greater psychological morbidity.

Analysis

We compared responses to simple and complex questions about anxiety and low mood to test the proposition that older people were more likely to endorse simple questions, which require less cognitive processing, than complex ones. The closest matches were as follows. The CIDI screen question for generalized anxiety disorder, 'In the past 12 months, have you had a period of a month or more when for most of the time you felt worried, tense or anxious about everyday problems such as work or family?', was contrasted with the K-10 item, 'In the past four weeks, about how often did you feel nervous?' The CIDI depression screen question, 'In the past 12 months, have you had two weeks or longer when nearly every day you felt sad, empty or depressed for most of the day?', was contrasted with its closest K-10 match, 'In the past four weeks, about how often did you feel depressed?'

Table 1. Cross-tabulation of numbers of positive and negative responses to (a) selected K-10 and CIDI anxiety questions and (b) selected K-10 and CIDI depression questions

(a)		CIDI anxiety question		
		Negative	Positive	Total
K-10 anxiety question	Negative	8284 ^a	2071 ^b	10 355
	Positive	83 ^c	203 ^d	286
	Total	8367	2274	10 641
(b)		CIDI depression question		
		Negative	Positive	Total
K-10 depression question	Negative	9121 ^a	1114 ^b	10 235
	Positive	131 ^c	275 ^d	406
	Total	9252	1389	10 641

Superscript letters a–d denote table cells.

Logically, a person who endorses a K-10 symptom as present 'all or most of the time' in the previous 4 weeks will also endorse the related CIDI item that refers to 'most of the time' for a month or more within the past 12 months. However, the interview takes 1 to 2 hours to administer and covers multiple social, mental, physical and service domains so complete consistency is unlikely. We explore in this paper the relative inconsistency in responses to selected K-10 and CIDI items of younger and older subjects, using logistic regression analysis to control for covariates or their proxies that could plausibly impinge on cognitive processing skills.

Only the 286 respondents who responded positively to the K-10 anxiety question were considered in the analysis concerning anxiety and only the 406 respondents who responded positively to the K-10 depression question were considered in the analysis concerning depression. Patterns of response to the K-10 and CIDI anxiety and depression questions are shown in Table 1(a, b) respectively. Note that the CIDI questions concerned a 12-month period and therefore generated more positive responses.

Our analyses concern only respondents who responded 'yes' to K-10 (Table 1, cells c and d). Within these restricted classes, agreement (yes/no) with the related CIDI screen question was taken as the dependent variable. Covariates in the regression equation included age (18–44, 45–64, ≥65 years), gender, country of birth (English speaking; non-English speaking), educational qualification (none or basic; skilled or tertiary) and self-rated health (excellent to good; fair to poor). Covariates were entered simultaneously in the regression equation. MMSE scores




Fig. 1. The 12-month prevalence of CIDI-diagnosed ICD-10 anxiety and depressive disorders, and mean scores on K-10 and GHQ-12, as a percentage of the index age group 18–24 years.

could not be used as the test was not administered to all respondents.

Results

Mean K-10 scores fell from 15.2 (s.d. = 5.1) in the age group 18–24 years to 13.2 (s.d. = 3.9) in the ≥75-year band, a difference of 2.0 points out of a maximum of 50. Fig. 1 maps K-10 scores, GHQ-12 scores and ICD-10 disorders across the lifespan, using age band 18–24 years as the index group. For respondents aged ≥75 years, 12-month prevalence rates were 19.2% of those in the index group for ICD-10 anxiety disorders and




Fig. 2. Disagreement rates (%) between K-10 and CIDI anxiety and depression questions.

3.7% for major depression and dysthymia. K-10 showed much less change. For respondents aged ≥ 75 years, mean scores were 87.0% of those in the index group.

Of the 286 respondents who endorsed the K-10 item 'In the past four weeks, about how often did you feel nervous?' as 'all or most of the time', 83 denied the CIDI anxiety screen question 'In the past 12 months, have you had a period of a month or more when for most of the time you felt worried, tense or anxious about everyday problems such as work or family?' Disagreement rates (CIDI-/K10+) increased from 29% in the 55-64 years band to 71% in those aged ≥ 75 years (Fig. 2).

On logistic regression analysis, disagreement rates were significantly higher for the age group ≥ 65 years [odds ratio (OR) 1.61, 95% confidence interval (CI) 1.07-2.44, $p=0.02$] and for those in good health (OR 2.34, 95% CI 1.33-4.11, $p=0.003$). Gender, country of birth and education made no significant contribution.

Of the 406 respondents who endorsed the K-10 item 'In the past four weeks, about how often did you feel depressed?' as 'all or most of the time', 131 denied the CIDI depression screen item 'In the past 12 months, have you had two weeks or longer when nearly every day you felt sad, empty or depressed for most of the day?' Disagreement rates (CIDI-/K10+) increased from 21% in the 55-64 years band to 71% in those aged ≥ 75 years (Fig. 2). On logistic regression analysis, the findings for age were OR 1.50 (95% CI 1.09-2.05, $p=0.01$). Other variables played no significant role.

Discussion

The National Survey has met its objective of generating interest in the prevalence and consequences of mental disorder in Australia. Its findings are shaping public policy and it is important to subject them to critical analysis because errors of fact or interpretation could have onerous consequences.

Although a decline in mental disorder with age makes sense, its rate may be exaggerated by dint of the length and complexity of structured mental health surveys. It is possible that aged respondents reply 'no' to questions when they exceed a certain, individually determined level of complexity. We cannot prove this, of course, but our hypothesis is supported by the observation that older people were more likely than younger ones to give answers to simple questions that failed to match their answers to convoluted questions concerning anxiety and depression (this trend might have been greater still had not low scorers on the MMSE been excluded). It also helps to explain why scores on K-10 and GHQ-12 showed much less attenuation with age than CIDI-generated psychiatric diagnoses.

Additional support for our hypothesis comes from a small, carefully controlled experimental study in which 31 older respondents endorsed CIDI depressive symptoms just as frequently as 32 younger ones but were more likely to attribute their symptoms to physical illness, thus reducing the likelihood of qualifying for a depressive syndrome. On regression analysis, physical attribution was associated more with reduced working memory than with age or objective markers of physical ill-health (Knäuper & Wittchen, 1994). If this finding is upheld in larger naturalistic and experimental studies, and in various clinical and cultural contexts, it will be important to reflect on ways in which structured interviews can be made more 'age friendly'.

Our analysis is limited. K-10 and CIDI questions do not tally exactly, especially in time-frame, and our match of questions is approximate. Alternative explanations for the rise in inconsistency of responses with age include fatigue (the mental health scales were administered earlier in the interview), a cohort effect (e.g. lower levels of psychological introspection in earlier generations) and reduced salience. In addition, it is likely that some younger people (e.g. those with limited language skills) will find CIDI a greater challenge than older people do.

We had expected to find that education and language skills also contributed to rates of inconsistency but this proved not to be the case, perhaps through lack of statistical power. People who rated their health as 'good to excellent' were more likely to respond

differently to simple and complex questions about anxiety but not depression, perhaps because those in poor physical health are so chronically anxious that symptoms remain to the front of their minds, irrespective of the complexity of the questions put to them.

The fact that scores on K-10 and GHQ-12 show limited change with age sends a warning not to ignore the symptoms and disability experienced by older people as a result of anxiety and depression. Both scales have robust psychometric properties and show close convergence with DSM and ICD diagnoses in younger adult age groups. There is a good argument for including at least one such scale in epidemiological surveys.

Declaration of Interest

None.

References

- Anderson, E** (2008). Cognitive change in old age. In *Oxford Textbook of Old Age Psychiatry*, 4th edn (ed. R. Jacoby, C. Oppenheimer, T. Denning and A. Thomas), pp. 33–50. Oxford University Press: Oxford.
- Copeland J, Kelleher M, Kellet J, Gourlay AJ, Gurland BJ, Fleiss JL, Sharpe L** (1976). A semi-structured clinical interview for the assessment and diagnosis of mental state in the elderly: Geriatric Mental State Schedule. 1: Development and reliability. *Psychological Medicine* **6**, 439–449.
- ESEMEd/MHEDEA 2000 Investigators** (2004). Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMEd) project. *Acta Psychiatrica Scandinavica* **109** (Suppl. 420), 21–27.
- Folstein MF, Folstein SE, McHugh PR** (1975). Mini-mental state: a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research* **12**, 189–198.
- Goldberg DP** (1972). *The Detection of Psychiatric Illness by Questionnaire*. Oxford University Press: London.
- Henderson AS** (1994). Does ageing protect against depression? *Social Psychiatry and Psychiatric Epidemiology* **29**, 107–109.
- Henderson S, Andrews G, Hall W** (2000). Australia's mental health: an overview of the general population survey. *Australian and New Zealand Journal of Psychiatry* **34**, 197–205.
- Kay DWK, Henderson AS, Scott R, Wilson J, Rickwood D, Grayson DA** (1985). Dementia and depression among the elderly living in the Hobart community: the effect of the diagnostic criteria on the prevalence rates. *Psychological Medicine* **15**, 771–788.
- Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand SLT, Walters EE, Zaslavsky AM** (2002). Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychological Medicine* **32**, 959–976.
- Kessler RC, Berglund P, Demler O, Jin R, Walters EE** (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Journal of the American Medical Association* **293**, 593–602.
- Knäuper B, Wittchen HU** (1994). Diagnosing major depression in the elderly: evidence for response bias in standardised diagnostic interviews? *Journal of Psychiatric Research* **28**, 147–164.
- Lee S, Tsang A, Zhang M-Y, Huang YQ, He YL, Liu ZR, Shen YC, Kessler RC** (2007). Lifetime prevalence and inter-cohort variation in DSM-IV disorders in metropolitan China. *Psychological Medicine* **37**, 61–73.
- Lindesay J, Baillon S, Brugha T, Dennis M, Stewart R, Araya R, Meltzer H** (2006). Worry content across the lifespan: an analysis of 16- to 74-year old participants in the British National Survey of Psychiatric Morbidity 2000. *Psychological Medicine* **36**, 1625–1633.
- O'Connor DW** (2006). Do older Australians truly have low rates of anxiety and depression? A critique of the 1997 National Survey of Mental Health and Wellbeing. *Australian and New Zealand Journal of Psychiatry* **40**, 623–631.
- Robins LN, Helzer JE, Croughan J, Ratcliff KS** (1981). National Institute of Mental Health Diagnostic Interview Schedule: its history, characteristics, and validity. *Archives of General Psychiatry* **38**, 381–389.
- Roth M, Huppert FA, Tym E, Mountjoy CQ** (1988). *The Cambridge Examination for Mental Disorders of the Elderly: CAMDEX*. Cambridge University Press: Cambridge.
- Villamil E, Huppert FA, Melzer D** (2006). Low prevalence of depression and anxiety is linked to statutory retirement ages rather than personal work exit: a national survey. *Psychological Medicine* **36**, 999–1009.
- Wells JE, Oakley Browne MA, Scott KM, McGee MA, Baxter J, Kokaua J** (2006). Prevalence, interference with life and severity of 12 month DSM-IV disorders in the Te Rau Hinengaro: the New Zealand Mental Health Survey. *Australian and New Zealand Journal of Psychiatry* **40**, 845–854.
- WHO** (1994). *Composite International Diagnostic Interview (CIDI): Researcher's Manual*. World Health Organization: Geneva.
- Yang Y** (2007). Is old age depressing? Growth trajectories and cohort variations in late-life depression. *Journal of Health and Social Behavior* **48**, 16–32.